VOL. 652, NO. 1 JOURNAL OF CHROMATOGRAPHY A OCTOBER 15, 1993

CONTENTS

5TH INTERNATIONAL SYMPOSIUM ON HIGH PERFORMANCE CAPILLARY ELECTROPHORESIS, ORLANDO, FL, JANUARY 25–28, 1993

PART I: VOLUME 652, NO. 1

Foreword
by B.L. Karger (Boston, MA, USA) 1

DNA APPLICATIONS

Capillary electrophoresis of DNA in uncross-linked polymer solutions
by A.E. Barron, D.S. Soane and H.W. Blanch (Berkeley, CA, USA) 3

Polymeric separation media for electrophoresis: cross-linked systems or entangled solutions (Review)
by Y.C. Bae and D. Soane (Hayward, CA, USA) 17

Simultaneous determination of the migration coefficient of each base in heterogeneous oligo-DNA by gel filled capillary electrophoresis
by T. Satow and T. Akiyama (Tokyo, Japan), A. Machida (Sapporo, Japan) and Y. Utagawa and H. Kobayashi (Saitama-ken, Japan) 23

Movement of DNA fragments during capillary zone electrophoresis in liquid polyacrylamide
by M. Chiari, M. Nesi and P.G. Righetti (Milan, Italy) 31

Use of glucomannan for the separation of DNA fragments by capillary electrophoresis
by T. Izumi, M. Yamaguchi, K. Yoneda, T. Isobe, T. Okuyama and T. Shinoda (Tokyo, Japan) 41

Separation of DNA restriction fragments by polymer-solution capillary zone electrophoresis. Influence of polymer concentration and ion-pairing reagents
by R.P. Singhal and J. Xian (Wichita, KS, USA) 47

Separation of DNA fragments by capillary electrophoresis using replaceable linear polyacrylamide matrices
by Y.F. Pariat, J. Berka, D.N. Heiger, T. Schmitt, M. Vilenchik, A.S. Cohen, F. Foret and B.L. Karger (Boston, MA, USA) 57

Optimization of capillary electrophoretic separation of DNA fragments based on polymer filled capillaries
by D.A. McGregor and E.S. Yeung (Ames, IA, USA) 67

Capillary electrophoresis of polymerase chain reaction-amplified DNA using fluorescence detection with an intercalating dye
by B.R. McCord,, D.L. McClure and J.M. Jung (Quantico, VA, USA) 75

Electrophoretic separations of polymerase chain reaction-amplified DNA fragments in DNA typing using a capillary electrophoresis–laser induced fluorescence system
by K. Srinivasan (Gaithersburg, MD, USA), J.E. Girard (Washington, DC, USA), P. Williams, R.K. Roby and V.W. Weedn (Washington, DC, USA), S.C. Morris (Columbia, MD, USA) and M.C. Kline and D.J. Reeder (Gaithersburg, MD, USA) 83

Effect of urea concentration on the base-specific separation of oligodeoxynucleotides in capillary affinity gel electrophoresis
by Y. Baba and M. Tsuhako (Kobe, Japan) and T. Sawa and M. Akashi (Kagoshima, Japan) 93

Capillary gel electrophoresis and the analysis of DNA phosphorothioates
by L. DeDionisio (Foster City, CA, USA) 101

PROTEIN APPLICATIONS

Utility of high resolution capillary electrophoresis for monitoring peptide homo- and hetero-dimer formation
by J.P. Landers, R.P. Oda, J.A. Liebenow and T.C. Spelsberg (Rochester, MN, USA) 109

Capillary electrophoresis of abnormal hemoglobins associated with α-thalassemias
by M. Zhu, T. Wehr, V. Levi, R. Rodriguez, K. Shiffer and Z.A. Cao (Hercules, CA, USA) 119

Effect of detergents on the electrophoretic behaviour of plasma apolipoproteins in capillary electrophoresis
by T. Tadey and W.C. Purdy (Montreal, Canada) 131

Evaluation of sodium dodecyl sulfate non-acrylamide, polymer gel-filled capillary electrophoresis for molecular size separation of recombinant bovine somatotropin
by K. Tsuji (Kalamazoo, MI, USA) 139

Characterization and performance of a neutral hydrophilic coating for the capillary electrophoretic separation of biopolymers
by D. Schmalzing, C.A. Piggee, F. Foret, E. Carrilho and B.L. Karger (Boston, MA, USA) 149

High-efficiency capillary electrophoretic separation of basic proteins using coated capillaries and cationic buffer additives. Evaluation of protein–capillary wall interactions
by A. Cifuentes, J.M. Santos, M. de Frutos and J.C. Diez-Masa (Madrid, Spain) 161

Buffer conditions affecting the separation of Maillard reaction products by capillary electrophoresis
by A.J. Tomlinson and J.P. Landers (Rochester, MN, USA), I.A.S. Lewis (London, UK) and S. Naylor (Rochester, MN, USA) 171

Capillary electrophoresis as a tool for the analysis of protein folding
by M.A. Strege and A.L. Lagu (Indianapolis, IN, USA) 179

High-performance capillary electrophoresis for the determination of trypsin and chymotrypsin inhibitors and their association with trypsin, chymotrypsin and monoclonal antibodies
by A.M. Arentoft (Frederiksberg, Denmark), H. Frøkiær (Lyngby, Denmark), S. Michaelsen (Tjele, Denmark), H. Sørensen (Frederiksberg, Denmark) and S. Sørensen (Lyngby, Denmark) 189

Characterization by capillary electrophoresis of the surface glycoproteins of ovine lentiviruses before and after treatment with glycosidic enzymes
by M.J. Schmerr and K.R. Goodwin (Ames, IA, USA) 199

Determination of milk proteins by capillary electrophoresis
by N. de Jong, S. Visser and C. Olieman (Ede, Netherlands) 207

Analysis of a recombinant granulocyte macrophage colony stimulating factor dosage form by capillary electrophoresis, capillary isoelectric focusing and high-performance liquid chromatography
by G.G. Yowell, S.D. Fazio and R.V. Vivilecchia (East Hanover, NJ, USA) 215

Separation of amino acid–oxazole derivatives of the redox coenzyme pyrroloquinoline quinone by capillary zone electrophoresis
by Y. Esaka, Y. Yamaguchi, K. Kano and M. Goto (Gifu, Japan) 225

CHIRAL SEPARATIONS

Capillary electrophoretic chiral separations using β-cyclodextrin as resolving agent. II. Bases: chiral selectivity as a function of pH and the concentration of β-cyclodextrin
by Y.Y. Rawjee, R.L. Williams and Gy. Vigh (College Station, TX, USA) 233

Chiral separations using an immobilized protein–dextran polymer network in affinity capillary electrophoresis
by P. Sun, G.E. Barker and R.A. Hartwick (Binghamton, NY, USA), N. Grinberg (Rahway, NJ, USA) and R. Kaliszan (Gdansk, Poland) 247

Optical resolution by high-performance capillary electrophoresis. Micellar electrokinetic chromatography with sodium N-dodecanoyl-L-glutamate and digitonin
by K. Otsuka, M. Kashihara, Y. Kawaguchi, R. Koike and T. Hisamitsu (Osaka, Japan) and S. Terabe (Hyogo, Japan) 253

Separation of positional isomers and enantiomers using capillary zone electrophoresis with neutral and charged cyclodextrins
by N.W. Smith (Greenford, UK) 259

FUNDAMENTALS

Timescales of transient processes in capillary electrophoresis
by E.V. Dose and G. Guiochon (Knoxville and Oak Ridge, TN, USA) 263

Leakage current consideration of capillary electrophoresis under electroosmotic control
by C.-T. Wu, T.-L. Huang and C.S. Lee (Baltimore, MD, USA) 277

Dispersion in capillary electrophoresis with external flow control methods
by C.A. Keely, R.R. Holloway, T.A.A.M. van de Goor and D. McManigill (Palo Alto, CA, USA) 283

On-line measurement of electroosmosis in capillary electrophoresis using a conductivity cell
by B.J. Wanders, T.A.A.M. van de Goor and F.M. Everaerts (Eindhoven, Netherlands) 291

In vitro observation of interactions of iron and transferrin by capillary isoelectric focusing with a concentration gradient imaging detection system
by J. Wu and J. Pawliszyn (Waterloo, Canada) 295

PART II: VOLUME 652, NO. 2

Automated pK_a determination at low solute concentrations by capillary electrophoresis
by J.A. Cleveland, Jr. and M.H. Benko (Indianapolis, IN, USA) and S.J. Gluck and Y.M. Walbroehl (Midland, MI, USA) 301

Characterization of quartz capillaries for capillary electrophoresis
by J. Kohr and H. Engelhardt (Saarbrücken, Germany) 309

Thermally induced fluctuations of the electric current and baseline in capillary electrophoresis
by M.S. Bello, P. de Besi and P.G. Righetti (Milan, Italy) 317

Computer-assisted determination of the inner temperature and peak correction for capillary electrophoresis
by M.S. Bello, E.I. Levin and P.G. Righetti (Milan, Italy) 329

Computer-assisted pH optimization for the separation of geometric isomers in capillary zone electrophoresis
by J.C. Jacquier, C. Rony and P.L. Desbene (Evreux and Mont Saint Aignan, France) 337

DETECTION

High-efficiency filter fluorometer for capillary electrophoresis and its application to fluorescein thiocarbamyl amino acids
by E. Arriaga, D.Y. Chen, X.L. Cheng and N.J. Dovichi (Edmonton, Canada) 347

Semiconductor laser-induced fluorescence detection in capillary electrophoresis using a cyanine dye
by F.-T.A. Chen, A. Tusak, S. Pentoney, Jr., K. Konrad, C. Lew, E. Koh and J. Sternberg (Fullerton, CA, USA) 355

On-line chemiluminescence detection of proteins separated by capillary zone electrophoresis
by T. Hara, J. Yokogi, S. Okamura, S. Kato and R. Nakajima (Kyoto, Japan) 361

Cetyltrimethylammonium chloride as a surfactant buffer additive for reversed-polarity capillary electrophoresis–electrospray mass spectrometry
by J. Varghese and R.B. Cole (New Orleans, LA, USA) 369

Determination of peptides by capillary electrophoresis–electrochemical detection using on-column Cu(II) complexation
by M. Deacon, T.J. O'Shea and S.M. Lunte (Lawrence, KS, USA) and M.R. Smyth (Dublin, Ireland) 377

Capillary electrophoresis of inorganic cations and low-molecular-mass amines using a copper-based electrolyte with indirect UV detection
by J.M. Riviello and M.P. Harrold (Sunnyvale, CA, USA) 385

CLINICAL/PHARMACEUTICAL APPLICATIONS

In vivo monitoring of glutamate in the brain by microdialysis and capillary electrophoresis with laser-induced fluorescence detection
by L. Hernandez, S. Tucci, N. Guzman and X. Paez (Merida, Venezuela) 393

Collinear laser-induced fluorescence detector for capillary electrophoresis. Analysis of glutamic acid in brain dialysates
by L. Hernandez (Merida, Venezuela and Ramonville, France), N. Joshi and E. Murzi (Merida, Venezuela), P. Verdeguer and J.C. Mifsud (Ramonville, France) and N. Guzman (Princeton, NJ, USA) 399

Assessment of automated capillary electrophoresis for therapeutic and diagnostic drug monitoring: determination of bupivacaine in drain fluid and antipyrine in plasma
by H. Wolfisberg, A. Schmutz, R. Stotzer and W. Thormann (Berne, Switzerland) 407

Investigation of the metabolism of the neuroleptic drug haloperidol by capillary electrophoresis
by A.J. Tomlinson, L.M. Benson, J.P. Landers and G.F. Scanlan (Rochester, MN, USA), J. Fang (Saskatoon, Canada), J.W. Gorrod (London, UK) and S. Naylor (Rochester, MN, USA) 417

Separation of estrogens and rodenticides using capillary electrophoresis with aqueous–methanolic buffers (Short Communication)
by K.J. Potter, R.J.B. Allington and J. Algaier (Lincoln, NE, USA) 427

Separation of gangliosides using cyclodextrin in capillary zone electrophoresis
by Y.S. Yoo, Y.S. Kim, G.-J. Jhon and J. Park (Seoul, South Korea) 431

Screening for diuretics in urine and blood serum by capillary zone electrophoresis
by J. Jumppanen, H. Sirén and M.-L. Riekkola (Helsinki, Finland) 441

Effect of buffer solution pH on the elution and separation of β-blockers by micellar electrokinetic capillary chromatography
by P. Lukkari, H. Vuorela and M.-L. Riekkola (Helsinki, Finland) 451

Validation of a capillary electrophoresis method for the determination of a quinolone antibiotic and its related impurities
by K.D. Altria and Y.L. Chanter (Ware, UK) 459

Sample matrix effects in capillary electrophoresis. I. Basic considerations
by L.L. Garcia and Z.K. Shihabi (Winston-Salem, NC, USA) 465

Sample matrix effects in capillary electrophoresis. II. Acetonitrile deproteinization
by Z.K. Shihabi (Winston-Salem, NC, USA) 471

Determination of flavonoids by micellar electrokinetic capillary chromatography
by C. Bjergegaard, S. Michaelsen, K. Mortensen and H. Sørensen (Frederiksberg, Denmark) 477

Determination of theophylline in plasma using different capillary electrophoretic systems
by I.M. Johansson, M.-B. Grön-Rydberg and B. Schmekel (Uppsala, Sweden) 487

Separation of water-soluble vitamins via high-performance capillary electrophoresis
by U. Jegle (Waldbronn, Germany) 495

MISCELLANEOUS APPLICATIONS

Separation and determination of glycosaminoglycan disaccharides by micellar electrokinetic capillary chromatography for studies of pelt glycosaminoglycans
by S. Michaelsen (Tjele, Denmark) and M.-B. Schrøder and H. Sørensen (Frederiksberg, Denmark) 503

Determination of oligosaccharides by capillary zone electrophoresis
by A.M. Arentoft, S. Michaelsen and H. Sørensen (Frederiksberg, Denmark) 517

Micellar electrokinetic capillary chromatography analysis of the behavior of bilirubin in micellar solutions
by A.D. Harman, R.G. Kibbey, M.A. Sablik, Y. Fintschenko, W.E. Kurtin and M.M. Bushey (San Antonio, TX, USA) 525

Isolation and quantification of ergovaline from *Festuca arundinacea* (tall fescue) infected with the fungus *Acremonium coenophialum* by high-performance capillary electrophoresis
by Y. Ma, K.G. Meyer and D. Afzal (Kirksville, MO, USA) and E.A. Agena (Indianola, IA, USA) 535

Ion-association capillary electrophoresis. New separation mode for equally and highly charged metal chelates
by N. Iki, H. Hoshino and T. Yotsuyanagi (Sendai, Japan) 539

Determination of organolead and organoselenium compounds by micellar electrokinetic chromatography
by C.L. Ng, H.K. Lee and S.F.Y. Li (Singapore, Singapore) 547

Use of capillary electrophoresis for monitoring citrus juice composition
by P.F. Cancalon and C.R. Bryan (Lake Alfred, FL, USA) 555

Quantitation of organic acids in sugar refinery juices with capillary zone electrophoresis and indirect UV detection
by S.P.D. Lalljie, J. Vindevogel and P. Sandra (Ghent, Belgium) 563

AUTHOR INDEX 571